Groupe d'Annecy

Laboratoire d'Annecy-le-Vieux de Physique des Particules

ENSLAPP

Groupe de Lyon

Ecole Normale Supérieure de Lyon

arXiv:hep-ph/9712342v1 11 Dec 1997

The Standard Model of particle physics*

P. Aurenche[1]

Laboratoire de Physique Théorique ENSLAPP†

[1] Groupe d'Annecy: LAPP, BP 110, F-74941 Annecy-le-Vieux Cedex, France.

Abstract

A brief review of the Standard Model of particle physics is presented.

hep-ph/9712342
ENSLAPP-A-659/97
July 1997

*Lectures presented at the Vth Séminaire Rhôdanien de Physique, Dolomieu, March 1997

†URA 14-36 du CNRS, associée à l'Ecole Normale Supérieure de Lyon et à l'Université de Savoie.

1 Introduction

From the experimental point of view the world of "elementary particles" consists in:
the leptons: they have spin $\frac{1}{2}$ and come in three doublets, (e^-, ν_e) the electron and its associated neutrino, (μ^-, ν_μ) the muon and its neutrino, (τ^-, ν_τ) the tau and its neutrino.
the vector bosons: they have spin 1 and there is a massless boson, the photon, and three massive ones, the W^+, W^- and the Z.
the hadrons: one distinguishes the mesons of integer spin ($S = 0, 1, \ldots$) from the baryons of half-integer spin ($S = \frac{1}{2}, \frac{3}{2}, \ \ldots$).
The *leptons* and the *vector bosons* are structureless down to a scale of about 10^{-3} to 10^{-4} fermi, *i.e.* 10^{-18} to 10^{-19} m according to the most recent experimental results and they are treated as elementary fields appearing in lagrangians which describe the dynamics of their interactions. One the other hand, the *hadrons* have been known for a long time to have a finite size (typically of the order of 1 fm) and there exist so many hadrons (about 150 mesons and 120 baryons) that they cannot be considered as elementary.
Three types of forces have been identified which describe the dynamics of particles: the strong force which affects only the hadrons, and the electromagnetic and weak forces.

The basic principle which guides the construction of models of particle physics is that of local gauge invariance according to which the physical properties do not depend on the phases of the fields. The Standard Model is a (highly successful) example of a minimal model based on the local gauge group

$$SU(3) \otimes SU(2)_L \otimes U(1)_Y$$

i.e. the direct product of three simple groups. The dynamical consequences of this local gauge invariance are presented in more details in the next sections. We just state now the main features of each of these groups.

– The $SU(3)$ gauge group or colour group is the symmetry group of strong interactions. This group acts on the quarks which are the elementary constituents of matter and the interaction force is mediated by the gluons which are the gauge bosons of the group. The quarks and the gluons are coloured fields. The "coupling" (fine structure constant) between quarks and gluons is denoted by α_s which can be ≥ 1. Under some conditions, however, α_s is very small and perturbation theory applies. The $SU(3)$ colour symmetry is exact and consequently the gluons are massless. The theory of strong interactions based on colour $SU(3)$ is called Quantum Chromodynamics.

– The $SU(2)_L \otimes U(1)_Y$ is the gauge group of the unified weak and electromagnetic interactions, where $SU(2)_L$ is the weak isospin group, acting on left-handed fermions, and $U(1)_Y$ is the hypercharge group. At "low" energy ($M < 250$ GeV) the $SU(2)_L \otimes U(1)_Y$ symmetry is "spontaneously" broken and the residual group is $U(1)_{emg}$ whose generator is a linear combination of the $U(1)_Y$ generator and a generator of $SU(2)_L$: the corresponding gauge boson is of course the photon and the associated "coupling" is $\alpha \simeq \frac{1}{137}$. Symmetry breaking implies that the other gauge bosons acquire a mass: they are the heavy $W^\pm, Z$ bosons dicovered at CERN in the mid '80's. The symmetry breaking mechanism is associated to the names of Brout, Englert, Higgs and Kibble, but, for some reasons, it is often simply referred to as the Higgs mechanism. The electro-weak theory, based on spontaneously broken $SU(2)_L \otimes U(1)_Y$

gauge invariance, is knowned as the Glashow-Salam-Weinberg (GSW) model.

We discuss first the hadronic world and Quantum Chromodynamics (QCD). In a second part, we study the electro-weak force and the Glashow-Salam-Weinberg model.

2 Quantum Chromodynamics

The quark model was introduced more than 30 years ago to describe the hadrons and we know now that six quarks are necessary, *i.e.* the quarks come in six flavours denoted $q_a = u, d, s, c, b, t$ for quark up, down, strange, charm, bottom and top. Like the leptons, the quarks are point-like, $spin = \frac{1}{2}$ fields so that assuming a baryon is made up of 3 quarks, $B = (q_a q_b q_c)$, and a meson is made up of a quark-antiquark pair, $M = (q_a \overline{q_b})$, one obtains integer spins for the mesons and half-integer spin for baryons if account is taken of the relative orbital angular momentum of the quarks in the hadrons. The quarks d, s, b have charge $-\frac{1}{3}$ and the quarks u, c, t have charge $\frac{2}{3}$ so that hadrons have only integral charges. The problem is that a good description of the mass spectrum is obtained in the quark model with a baryonic wave function which is symmetrical under interchange of, *e.g.*, $q_a \leftrightarrow q_b$: this violates Fermi statistics. This problem can be cured if we introduce a new quantum number, the colour, and assume that each quark flavour comes in three colours:

$$(q_a)^T = (q_a^1, q_a^2, q_a^3) \quad \text{with } 1, 2, 3 \text{ the colour indices.} \tag{1}$$

The baryon wave function is then assumed to be totally antisymmetric under colour interchange,

$$B = \sum_{ijk} (q_a^i q_b^j q_c^k), \tag{2}$$

and the meson wave function is constructed as

$$M = \sum_i (q_a^i \overline{q_b^i}). \tag{3}$$

One introduces a group of transformation in colour space, $(q_a) \rightarrow (q'_a) = \mathcal{U}(q_a)$ where (q_a) is written as a column vector and $\mathcal{U}$ is a 3×3 matrix. In agreement with all experimental constraints, the group is chosen to be $SU(3)$, so that hadronic wave functions are colour singlets ($\mathbf{1}$), *i.e.* they are invariant under the action of the group. It is postulated that only colour $\mathbf{1}$ states are observable which, in agreement with data, excludes physical states of type $(q_a q_b)$ or $(q_a q_b \overline{q}_c)$ (this rules out colour groups such as $O(3)$ or $SO(3)$ for which such unobserved states can also be $\mathbf{1}$, but does not rule out $U(3)$ which will be discarded below on dynamical grounds). The colour hypothesis is further supported by the data on the rate of decay of $\pi^0 \rightarrow \gamma\gamma$ (via a quark loop) as well as the rate of hadron production in the high energy e^+e^- annihilation reactions (via the process $e^+e^- \rightarrow q_a^i \overline{q_a^i}$).

From now on we will take the quarks as the basic degrees of freedom of the hadronic world. Since quarks are confined into hadrons (no free quark observed!) one needs a dynamical mechanism to glue the quarks together. To introduce such a force one proceeds by analogy with Quantum Electrodynamics (QED) and one makes the colour symmetry a local gauge

symmetry. To enforce the invariance of the Lagrangian under a local $SU(3)$ transformation, it is necessary to introduce eight coloured vector fields which are in QCD the analogous of the photon in QED. In more details, the free fermion Lagrangian for one quark field of a given flavour is

$$\mathcal{L} = \bar{\psi}(i\ \partial\!\!\!/ - m)\psi = \bar{\psi}^1(i\ \partial\!\!\!/ - m)\psi^1 + \bar{\psi}^2(i\ \partial\!\!\!/ - m)\psi^2 + \bar{\psi}^3(i\ \partial\!\!\!/ - m)\psi^3$$

where ψ is a column vector, each coloured component of which is a Dirac spinor, $\psi^T = (\psi^{1^T}, \psi^{2^T}, \psi^{3^T})$, ψ belonging to the **3** (triplet) representation of SU(3). A local gauge transformation acting on the fundamental **3** representation is parameterised by

$$\mathcal{U} = e^{ig\sum_a \alpha_a(x)T_a},$$

with g the coupling. The T_a's are the eight 3×3 matrices generators of the group, and the $\alpha_a(x)$'s are eight arbitrary real parameters depending on space-time coordinates. The T_a's are traceless, hermitian and satisfy the commutation relations $[T_a, T_b] = if_{abc}T_c$, with f_{abc} totally antisymetric and real. A gauge transformation acting on ψ amounts to a locally dependent change of phase and for an infinitesimal transformation one has

$$\delta\psi = ig\alpha_a T^a\psi, \qquad \delta\partial_\mu\psi = ig\alpha_a T^a\partial_\mu\psi + ig(\partial_\mu\alpha_a)T^a\psi,$$

The Lagrangian is not gauge invariant since its variation is

$$\delta\mathcal{L} = \bar{\psi}\left\{-g(\partial_\mu\alpha_a)T^a\gamma^\mu\right\}\psi. \tag{4}$$

To regain invariance, it is sufficient to introduce 8 vector fields $A^b_\mu(x)$ transforming as

$$\delta A^c_\mu = \partial_\mu\alpha^c - f^{cab}\alpha^a A^b_\mu.$$

and consider the lagrangian density

$$\mathcal{L} = \bar{\psi}\left(i(\partial_\mu - igA^b_\mu(x)T^b)\gamma^\mu - m\right)\psi, \qquad b = 1, \ldots, 8. \tag{5}$$

To make the "gauge bosons" dynamical degrees of freedom one constructs the anti-symmetric stress-energy tensor

$$F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + gf^{abc}A^b_\mu A^c_\nu \tag{6}$$

and by analogy with QED the "gauge boson" kinetic term is proportional to the gauge invariant quantity $F^a_{\mu\nu}F^{a\mu\nu}$. The QCD lagrangian density takes then the form

$$\mathcal{L}_{\rm QCD} = -\frac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} \ + \ \bar{\psi}\left(i(\partial_\mu - igA^a_\mu T^a)\gamma^\mu - m\right)\psi \tag{7}$$

where summation over repeated indices is assumed. Gauge invariance excludes mass terms of the form $m_A^2 A^a_\mu A^{a\mu}$ so that the gauge bosons are massless and induce long distance forces. We remark at this point that if U(3) had been chosen as the gauge group one would have a ninth massless gauge boson, which would be coulourless (associated to the $\mathbf{1}$ generator). This would induce long distance strong interactions between hadrons in contradiction with

experiments which tell us that the nuclear force is short-range $\sim \frac{1}{m_\pi}$. A crucial difference between Chromodynamics and Electrodynamics is the existence of self-couplings of the gluons. This is a consequence of the non-abelian nature of the colour group which imposes the term linear in f^{abc} in eq. (6) and leads to three-gluon and four-gluon couplings, proportional to g and g^2 respectively, via the kinetic term $F^a_{\mu\nu}F^{a\mu\nu}$ in eq. (7). The consequences of this fact are crucial and justify the use of the perturbative approach to study hadronic interactions at high energies as we are now going to discuss qualitatively.

When calculating higher order diagrams in field theories like QED or QCD one encounters "ultraviolet" divergences, *i.e.* divergences at very high energies or equivalently at very short distances, related to the point-like structure of fields and the local nature of interactions. To give a meaning to the perturbative expansion one needs to carry out the procedure of "renormalisation" and redefine (renormalize) all the parameters appearing in the "bare" Lagrangian eq. (7) by fixing their values at some arbitrary point in momentum space. Expressed in terms of the renormalized parameters the perturbative series has finite coefficients. Applied to quark-quark scattering, the matrix element-squared, *i.e.* up to a flux factor the differential cross section, in the one-loop approximation (see figure)



keeps essentially the same form as at the zeroth order approximation namely

$$|\mathcal{M}^{q_1q_2\to q_1'q_2'}|^2 \equiv \alpha_s(q^2)|\bar{u}(q_2')\gamma_\mu u(q_2)\frac{1}{q^2}\bar{u}(q_1')\gamma^\mu u(q_1)|^2, \tag{8}$$

the main difference with the lowest order result being that the coupling constant, $\alpha_s(q^2) = g^2(q^2)/4\pi$, is running in the sense that it depends on the momentum scale q^2 charateristic of the process under consideration. More precisely, the calculation leads to

$$\alpha_s(q^2) = \frac{\alpha_s(\mu_0^2)}{1 + \frac{(11N_c - 2N_F)}{12\pi}\ \alpha_s(\mu_0^2)\ \ln(\frac{-q^2}{\mu_0^2})} \tag{9}$$

where $\alpha_s(\mu_0^2)$ is the renormalised coupling at an arbitrary energy scale μ_0 and q^2 is the momentum transfer of the scattering process. $N_c = 3$ is the number of colours and $N_{_F}$ is the number of active flavours . Since for QCD the combination $(11N_c - 2N_{_F})$ is positive the theory predicts that for large momentum transfers and therefore large energy the effective QCD coupling decreases and vanishes at asymptotically high energies: this is called asymptotic freedom. The smallness of the strong interaction coupling constant at high energies justifies the use of perturbation theory in the study of particle physics processes in particular at the colliders such as those of FERMILAB (proton-antiproton collisions at 1.8 TeV), HERA

(electron-proton collisions at about 300 GeV) or for hadronic proccesses in e^+e^- collisions at LEP from 90 GeV to 190 GeV. On the other hand, at very low energy, the strong interaction coupling is increasing as the energy decreases and becomes so large as to hopefully lead to the confinement of quarks and gluons into hadrons at the scale of roughly 1 GeV. Of course, perturbation theory is not applicable then and this domain is studied using lattice gauge methods. The non-abelian nature of QCD is crucial to obtain aymptotic freedom as for an abelian theory one would have $N_c = 0$ and the effective coupling would be rising with energy exactly as it is the case for QED.

The property of asymptotic freedom is not sufficient in itself for perturbation theory to be useful in the study of hadronic collisions at high energies. A crucial property of QCD is contained in the "factorisation theorem" which states that the interaction between hadrons at high energies is factorised into a non-perturbative part which contains the information about the distribution of partons (quarks and gluons) into the hadrons and a perturbative, calculable part which describes the interaction between the partons at short distance. This is illustrated in the following picture



and made quantitative in the equation giving the generic form of a scattering cross section between two hadrons

$$d\sigma^{H_1H_2} = \sum_{i,j} \int dx_1\ dx_2\ F_i^{H_1}(x_1, M)\ F_j^{H_2}(x_2, M) \left[\hat{\sigma}_{ij} + \alpha_s(M)\ \hat{\sigma}_{ij}^{(1)} + \alpha_s^2(M)\hat{\sigma}_{ij}^{(2)}...\right]. \quad (10)$$

The structure function (or parton density) $F_i^H(x)$ is related to the probability of finding, in hadron H, a parton of type i carrying a fraction x of the hadron momentum. The x dependence of $F_i^H(x)$ is not calculable in perturbative QCD. The series in square brackets is the cross section for parton-parton scattering at the energy scale M (or distance $1/M$) charateristic of the process under consideration. It is the part of the cross section which is calculable provided M is large enough so that $\alpha_s(M)$ is small. The theory also predicts the scale M dependence (scaling violations) of the structure functions.
Such cross section have been calculated for most relevant processes up to the second term in the series and sometimes up to the third term. A sustained theoretical effort is going on to resum to all orders large contributions which may appear in the coefficients $\hat{\sigma}_{ij}^{(n)}$.

Experimentally there has been for a long time evidence for gluons. Already in the early deep-inelastic experiments, where one studies the scattering of electrons off protons at large

momentum transfers, it was possible to measure the amount of momentum carried by the charged constituents of the proton (since the virtual photon exchanged between the electron and the proton couples only to charged fields). It was found that charged constituents (the quarks) carry only about half of the momentum of the proton, the rest being carried by neutral fields (the gluons?) assumed to bind the quarks together into the hadron. Later on, with the advent of very high energy colliders it became possible to "visualise" quarks and gluons as highly concentrated deposits of energy, called "jets" in hadronic calorimeters: these jets fit with the interpretation of being the hadronic decays of quarks and gluons produced at large momentum transfers. Perturbative Quantum Chromodynamics has been tested on many different observables, the most spectacular of which is probably the jet production cross section as a function of the jet transverse momentum in proton-antiproton collisions where experiment and theory agree over ten orders of magnitude.

For further details one may consult refs. [1]-[6]. For an up-to-date status of QCD phenomenological studies see ref. [7].

3 The Glashow-Salam-Weinberg model

Before entering the description of the unified theory of electro-weak interactions, based on broken gauge invariance, it is useful to briefly review the Fermi theory of weak interactions and its phenomenological extensions: this will serve to motivate the choice of the gauge group $SU(2)_L \otimes U(1)_Y$ as well as illustrate the problems related to the presence of massive gauge bosons.

3.1 The Fermi theory and its extensions

At the beginning was the Fermi theory of neutron decay

$$n \to p\ e^-\ \overline{\nu}_e$$

(equivalently in the quark model, d quark decay $d \to u\ e^-\ \overline{\nu}_e$) and muon decay

$$\mu^- \to e^-\ \overline{\nu}_e\ \nu_\mu.$$

These transitions were described by a local current-current (4 fermion) interaction parameterised by the Lagrangian

$$\mathcal{L} = \frac{G}{\sqrt{2}} J^\dagger_\alpha(x) J^\alpha(x).$$

The current has a leptonic part and a hadronic part, $J_\alpha(x) = l_\alpha(x) + h_\alpha(x)$,

$$l_\alpha(x) = \bar{\psi}_e \gamma_\alpha(1-\gamma_5)\psi_{\nu e} + \bar{\psi}_\mu \gamma_\alpha(1-\gamma_5)\psi_{\nu\mu} + \cdots \quad (11)$$

$$h_\alpha(x) = \bar{\psi}_u \gamma_\alpha(1-\gamma_5)\psi_d + \bar{\psi}_c \gamma_\alpha(1-\gamma_5)\psi_s + \cdots. \quad (12)$$

The particular $V-A$ (vector$-$axial) form of the current, $\gamma_\alpha(1-\gamma_5)$, is dictated by experiment, in particular the angular distribution of the decay products. It means that (for massless fields)

only left-handed quarks or leptons $\psi_{_L} = (1-\gamma_5)\psi/2$ are sensitive to the weak interactions. The Fermi constant G is universal, *i.e.* it is the same for the hadronic sector and the leptonic sector and its value has been measured to be

$$G = 1.6639(2)10^{-5}\ \mathrm{GeV}^{-2}. \tag{13}$$

The transition matrix element for μ decay is

$$\mathcal{M} = \frac{G}{\sqrt{2}}(\bar{\psi}_e\gamma_\alpha(1-\gamma_5)\psi_{\nu_e})(\bar{\psi}_\mu\gamma^\alpha(1-\gamma_5)\psi_{\nu_\mu})^\dagger \tag{14}$$

which also describes, by crossing symmetry, the scattering process

$$\mu^-\ \nu_e \to e^-\ \nu_\mu \tag{15}$$

at the invariant energy squared $s = (p_{\nu_e} + p_\mu)^2$. The total cross section for the latter process is then found to be $\sigma \sim G^2 s$. This result could have easily been guessed on dimensional grounds since s is the only relevant scale of the problem besides the Fermi coupling G. However such a rapid rise of the cross section with energy cannot be asymptotically true as it violates the famous Froissart-Martin unitarity bound which requires $\sigma \leq \ln^2 s$ as $s \to \infty$. The problem is related to the locality of the current-current interaction. To improve the situation one introduces a massive charged particle to mediate the interaction between the two left-handed currents. It must be a vector particle because of the γ_α coupling in eq. (12). Denoting $M_{_W}$ the mass of this particle and $g_{_W}$ its dimensionless coupling to the currents, the matrix element eq. (14) becomes

$$\mathcal{M} = g_{_W}^2(\bar{\psi}_e\gamma_\alpha(1-\gamma_5)\psi_{\nu_e})\frac{g^{\alpha\beta} - \frac{q^\alpha q^\beta}{M_{_W}^2}}{q^2 - M_{_W}^2}(\bar{\psi}_\mu\gamma_\beta(1-\gamma_5)\psi_{\nu_\mu})^\dagger \tag{16}$$

where q^2 is the momentum transfer of the reaction. At low q^2 or s ($s \ll M_{_W}^2$) one exactly recovers the Fermi model provided one chooses

$$M_{_W}^2 = \sqrt{2}\ \frac{g_{_W}^2}{G} \tag{17}$$

while at high energies the total cross section is well behaved since it is easily derived that

$$\sigma \sim \frac{g^4}{M_{_W}^2}\frac{s}{s + M_{_W}^2}, \qquad s \gg M_{_W}^2. \tag{18}$$

It is useful at this point to comment on the differences between massless and massive vector bosons concerning the polarisation states. It is well known that a massless vector particle of momentum $q_\mu = (q_0, 0, 0, |q_0|)$, say, has two transverse polarisation (states) vectors which can be parameterised as

$$\varepsilon_{_T}^\mu = (0,1,0,0) \text{ and } (0,0,1,0) \tag{19}$$

whereas a massive vector particle of momentum $q_\mu = (q_0, 0, 0, \sqrt{q_0^2 - M_{_W}^2})$ has, besides the two transverse polarisations, a longitudinal one

$$\varepsilon_{_L}^{\mu} = \frac{1}{M_{_W}}(\sqrt{q_0^2 - M_{_W}^2}, 0, 0, q_0) \sim \frac{q_\mu}{M_{_W}} + \mathcal{O}(\frac{M_{_W}}{q_0}) \tag{20}$$

where the rightmost form is asymptotically valid ($|q_0| \gg M_{_W}$). If $W^\pm$ is a real particle it should be produced in reactions such as $e^+\ e^- \to W^+\ W^-$ or, less realistically, in $\nu_e\ \overline{\nu_e} \to W^+\ W^-$. Considering the latter reaction, the amplitude of which is given by only one Feynman diagram



Figure 1: *Possible Feynman diagrams for $\nu_e\bar{\nu}_e \to W^+W^-$ scattering. (***a***): e exchange; (***b***) hypothetical heavy electron E exchange; (***c***) neutral vector boson Z exchange.*

with the exchange of an electron (fig. 1a), we find the following results for the production of $W^\pm$ in transverse and longitudinal polarisation states respectively:

$$\begin{aligned} \sigma(\nu\bar{\nu} \to W_T^+ W_T^-) &\sim \text{constant} \\ \sigma(\nu\bar{\nu} \to W_L^+ W_L^-) &\sim \frac{g_{_W}^2}{M_{_W}^2}\ s \end{aligned} \tag{21}$$

The origin of the last expression can be understood if one recalls that the corresponding matrix element is asymptotically

$$\mathcal{M}_{\mu\nu}\ \varepsilon_{_L}^{\mu}(q)\varepsilon_{_L}^{\nu}(q') \sim \mathcal{M}_{\mu\nu}\frac{q_\mu}{M_{_W}}\frac{q'_\nu}{M_{_W}} \tag{22}$$

leading to the appearence of factors of s in the numerator. We thus find that the production of longitudinally polarised vector bosons violates the unitarity limit while that of transverse bosons is well behaved at high energies. Several ways have been tried to cure this problem: among them one can mention the hypothesis of a new heavy lepton (fig. 1b) and choose its couplings to enforce a proper behaviour of the cross section at high energies. It turns out that another possibility, namely that of a heavy neutral vector boson (fig. 1c), is realised in Nature. Assuming the most general couplings $(g_{_Z}\gamma_\mu(a - b\gamma_5),\ g'_{_Z}(p_{_{W^+}} + p_{_{W^-}})_\mu)$, they can be chosen to make cross sections such as $\nu\ e \to Z\ W$, $e^+\ e^- \to W^+\ W^-$, $\cdots$ asymptotically well behaved. However, this "bricolage" is not yet sufficient to have a satisfactory model. Indeed considering $e^+\ e^- \to Z\ Z$ scattering and keeping in principle small terms proportional to the mass of the electron, one finds an interference piece in the cross section $\sim g^4 m_e\sqrt{s}/M_{_Z}^4$ which again violates unitarity! Similar problems arise in the $W\ W$ scattering process, *e.g.* $W^+\ W^- \to W^+\ W^-$ which hopefully will be studied at LHC or in the future e^+e^- high energy

linear colliders: the cross sections for these processes diverge linearly in s. These problems can be solved by supposing the existence of a scalar particle which interacts with the bosons as well as the fermions with appropriately chosen couplings.

One can thus construct a viable electro-weak theory in the pedestrian way described above, carefully choosing masses and couplings of the newly introduced particles so as to ensure the correct behaviour of all cross sections. It is more instructive however to assume that these relations among masses and couplings arise from some hidden symmetry property. This is what we are going to do next. First we describe in some details the symmetry group assuming local gauge invariance. At this level, the chosen group requires all fields to be massless. The theory is renormalisable (well behaved at asymptotic energies) being a non-abelian field theory. Then, by the mechanism of "spontaneous symmetry breaking" whereby the symmetry of the lagrangian is preserved but the choice of a ground state breaks the symmetry, fermions and gauge bosons acquire a mass. After symmetry breaking, the theory remains renormalisable as a consequence of the underlying gauge invariance which imposes the required relations between couplings. One is left however with a large number of parameters (at least 19 for the Standard Model) which gives a motivation for a (still unsuccesful!) search of a deeper symmetry.

3.2 The $SU(2)_L$ gauge invariance

As discussed above, the weak interactions induce a transition between fermions of different charges. It is then natural to group fermions into doublets

$$\underbrace{\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}, \begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}, \begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}}_{\text{leptons}} ; \underbrace{\begin{pmatrix} u \\ d \end{pmatrix}, \begin{pmatrix} c \\ s \end{pmatrix}, \begin{pmatrix} t \\ b \end{pmatrix}}_{\text{quarks}} \tag{23}$$

Because of the $\gamma_\alpha(1-\gamma_5)$ form of the weak current, we have to distinguish between left-handed fermions which are sensitive to the weak interactions and the right handed part which are not. Therefore, each fermion field is decomposed as

$$\psi = \frac{1-\gamma_5}{2}\,\psi + \frac{1+\gamma_5}{2}\,\psi = \psi_{_L} + \psi_{_R}, \tag{24}$$

where $\psi_{_L}$ (resp. $\psi_{_R}$) is identified as the left (resp. right) component when $m_\psi = 0$. We introduce the left handed doublets:

$$\psi_{e_L} = \begin{pmatrix} \nu_e \\ e_{_L} \end{pmatrix}, \cdots \qquad \psi_{q_L} = \begin{pmatrix} u_{_L} \\ d_{_L} \end{pmatrix}, \cdots, \tag{25}$$

and the right-handed singlets $\psi_{e_R} = e_{_R}, \cdots, \psi_{q_R} = q_{_R}, \cdots$. The free massless fermion lagrangian is then written

$$\begin{aligned} -i\mathcal{L}_F &= \bar{\psi}_{e_{_L}} \not{\partial}\, \psi_{e_L} + \cdots + \bar{\psi}_{q_L}\, \not{\partial}\, \psi_{q_L} + \cdots + \bar{\psi}_{e_{_R}}\, \not{\partial}\, \psi_{e_R} + \cdots + \bar{\psi}_{q_R} \not{\partial}\, \psi_{q_R} + \cdots \\ &= \mathcal{L}_{F_L} + \mathcal{L}_{F_R}. \end{aligned} \tag{26}$$

We impose now a local SU(2) gauge invariance to the left handed part of the lagrangian, *i.e.* we impose $\delta\mathcal{L}_{F_L} = 0$ under the changes of phase

$$\delta\psi_{_L} = ig\left(\frac{\vec{\tau}}{2}\cdot\vec{\alpha}(x)\right)\psi_{_L} \quad , \quad \delta\bar{\psi}_{_L} = -ig\ \bar{\psi}_{_L}\left(\frac{\vec{\tau}}{2}\cdot\vec{\alpha}(x)\right) \tag{27}$$

where the 2×2 Pauli matrices $\vec{\tau} = (\tau_1, \tau_2, \tau_3)$ satisfy the algebra

$$\left[\frac{\tau_i}{2}, \frac{\tau_j}{2}\right] = i\ \epsilon_{ijk}\ \frac{\tau_k}{2}. \tag{28}$$

As for the SU(3) local invariance studied above, one has to introduce a multiplet of gauge vector fields $\vec{W}^{\mu}(x) = (W_1^{\mu}(x), W_2^{\mu}(x), W_3^{\mu}(x))$ transforming as

$$\delta\ W_i^{\mu}(x) = \partial^{\mu}\alpha_i(x) - \epsilon_{ijk}\alpha_i(x)\ W_k^{\mu}(x) \tag{29}$$

with $\vec{\alpha}(x) = (\alpha_1(x), \alpha_2(x), \alpha_3(x))$ three real parameters. The locally SU(2) gauge invariant left-handed lagrangian becomes then

$$\mathcal{L}_{F_L} = -\frac{1}{4}\ F_i^{\mu\nu}(x)F_{i_{\mu\nu}}(x) + \bar{\psi}_{_L} i\left(\partial^{\mu} - ig\frac{\vec{\tau}}{2}\cdot\vec{W}^{\mu}(x)\right)\gamma_{\mu}\psi_{_L} + \text{other fermions.} \tag{30}$$

where the antisymmetric tensor $F_i^{\mu\nu}(x)$ is

$$F_i^{\mu\nu}(x) = \partial^{\mu}W_i^{\nu}(x) - \partial^{\nu}W_i^{\mu}(x)\ + g\ \epsilon_{ijk}\ W_j^{\mu}(x)W_k^{\nu}(x). \tag{31}$$

(see eqs. (7), (6). We note at this point the perfect analogy between the construction of the "weak" lagrangian with that of QCD: the differences are in the choice of group which requires here only three vector bosons while for $SU(3)$ symmetry eight bosons had to be introduced. Also, the weak group acts only on the left handed components of the fields and consequently the $W_i^{\mu}(x)$ gauge bosons do not couple to the right handed fermion components. This is called the weak isospin group and is denoted $SU(2)_L$. The left handed fermion have isospin $t = 1/2$ and the third component (eigenvalue of $t_3 = \frac{\tau_3}{2}$) is $+1/2$ for $\nu_e, u_{_L}, ...$ and $-1/2$ for $e_{_L}, d_{_L}$. The right-handed components have of course isospin $t = 0$. We assume there are no right-handed neutrinos. Defining the charged fields $W^{\pm\mu}(x) = (W_1^{\mu} \mp iW_2^{\mu})/\sqrt{2}$ these two bosons will of course be identified with the charged bosons introduced in the phenomenological discussion above. One can read off their interaction term with the fermions from eq. (30) and one finds the familiar $(V-A)$ coupling $-i\frac{g}{\sqrt{2}}\bar{\nu}_e\gamma^{\mu}e_{_L}^{-}W_{\mu}^{+}$ and similarly for other fermion doublets (note the relation $g/2\sqrt{2} = g_{_W}$ of eq. (16)). However $W^{3\mu}$ cannot be the photon nor the Z boson since it also couples via $(V-A)$ in disagreement with data. Anyhow, since we need a fourth vector boson we introduce a new abelian local invariance and the point is to show that, by an astute choice of the associated charge, one can reconstruct the physical couplings. Before doing this let us remark the three and four boson self-interaction terms buried into the $F_i \cdot F_i$ term in the lagrangian density eq. (30) above.

3.3 The $U(1)_Y$ gauge invariance

The $SU(2)_L$ gauge invariance leads to a conserved current: for example for the electron sector it is

$$J_3^\mu(x) = \frac{1}{2}\bar{\psi}_{e_L}\gamma_\mu\tau_3\psi_{e_L} = \frac{1}{2}[\bar{\nu}\gamma_\mu\nu - \bar{e}_{_L}\gamma_\mu e_{_L}]. \tag{32}$$

On the other hand, we know that the electromagnetic current is conserved

$$J^\mu_{emg}(x) = e_{el}\ \bar{e}\gamma_\mu e = e_{el}\ [\bar{e}_{_L}\gamma_\mu e_{_L} + \bar{e}_{_R}\gamma_\mu e_{_R}], \tag{33}$$

where e_{el} is the electron charge expressed in units of the proton charge ($e_{el} = -1$). The combination $J^\mu_{emg}(x) - J^\mu_3(x)$ must then necessarily be conserved. Denoting it the hypercharge current we define

$$\frac{J^\mu_Y(x)}{2} \equiv J^\mu_{emg}(x) - J^\mu_3(x) = -\frac{1}{2}\ \bar{\psi}_{e_L}\gamma_\mu\psi_{e_L} - \bar{e}_{_R}\gamma_\mu e_{_R} \tag{34}$$

where we recall the definition of the spinor doublet ψ_{e_L} in eq. (25). Therefore the hypercharge $\frac{Y_{el}}{2} = e_{el} - (t_3)_{el}$ is conserved and its value can be immediately read off from eq. (34). One finds the following assignements: -1 for the left-handed doublet and -2 for the right-handed electron partner. The above relation is a particular example of the famous Gell-Mann/Nishijima formula

$$\frac{Y}{2} = Q - T_3 \tag{35}$$

which gives the hypercharge in terms of the electric charge and the isopin of the field. Associated to this conserved hypercharge we introduce a local abelian gauge symmetry denoted $U(1)_Y$ to which is associated a new gauge boson $B_\mu(x)$. The action of this transformation on the fermion fields is:

$$\delta\psi_{_L} = ig'\ y_{\psi_{_L}}\ \beta(x)\ \psi_{_L}, \qquad \delta\psi_{_R} = ig'\ y_{\psi_{_R}}\ \beta(x)\ \psi_{_R} \tag{36}$$

with g' the associated coupling constant and the hypercharge values of the $B_\mu(x)$ field couplings to the fermions are listed in the following table.

	t	t_3	Y	Q
ν_e	1/2	1/2	-1	0
$e_{_L}$	1/2	$-1/2$	-1	-1
$e_{_R}$	0	0	-2	-1
$u_{_L}$	1/2	1/2	1/3	2/3
$d_{_L}$	1/2	$-1/2$	1/3	$-1/3$
$u_{_R}$	0	0	4/3	2/3
$d_{_R}$	0	0	$-2/3$	$-1/3$

(37)

In summary, the initial free lagrangian eq. (26), becomes after imposing a $SU(2)$ local symmetry on the left-handed fields eq. (25) and an appropriate $U(1)$ invariance on both the

left-handed fields and a right-handed ones,

$$\begin{aligned}\mathcal{L} &= \mathcal{L}_G + \mathcal{L}_F \\ &= -\frac{1}{4}\, F_i^{\mu\nu}(x)\, F_i^{\mu\nu}(x) - \frac{1}{4}\, G^{\mu\nu}(x)\, G^{\mu\nu}(x) \\ &\quad + \bar{\psi}_{e_L}\, i\, \not{D}_{_L}\, \psi_{e_L} + \bar{\psi}_{q_L}\, i\, \not{D}_{_L}\, \psi_{q_L} + \\ &\quad + \bar{e}_{_R}\, i\, \not{D}_{_R}\, e_{_R} + \bar{u}_{_R}\, i\, \not{D}_{_R}\, u_{_R} + \bar{d}_{_R}\, i\, \not{D}_{_R}\, d_{_R}\end{aligned} \tag{38}$$

where only the electron and (u, d) quark families have been specified. The left covariant derivative acting on the field $\psi_{_L}$ is

$$D_L^\mu = \partial^\mu - i\, g\, \frac{\vec{\tau}}{2} \cdot \vec{W}^\mu - i\, g'\, y_{\psi_{_L}}\, B^\mu \tag{39}$$

and the right covariant derivative acting on field $\psi_{_R}$

$$D_R^\mu = \partial^\mu - i\, g'\, y_{\psi_{_R}}\, B^\mu. \tag{40}$$

The kinetic term of the gauge fields are constructed from

$$\begin{aligned} G^{\mu\nu}(x) &= \partial^\mu B^\nu(x) - \partial^\nu B^\mu(x) \qquad \text{(abelian field)} \\ F_i^{\mu\nu}(x) &= \partial^\mu W_i^\mu(x) - \partial^\nu W_i^\nu(x) + g\, \epsilon_{ijk}\, W_j^\mu(x)\, W_k^\nu(x). \end{aligned} \tag{41}$$

It is important to point out that $SU(2)_L \otimes U(1)_Y$ invariance imposes that all fermions are massless. Indeed a fermion mass term in the lagrangian would have the form

$$\mathcal{L}_{mass} = m\, \bar{\psi}\psi = m(\bar{\psi}_{_L}\psi_{_R} + \bar{\psi}_{_R}\psi_{_L}). \tag{42}$$

But since $\psi_{_L}$ is a doublet and $\bar{\psi}_{_R}$ a singlet under $SU(2)$, the mass term cannot be invariant under a gauge transformation!

It is clear from eq. (34) that the physical fields of the photon will be a linear combination of W_μ^3 and the gauge boson B_μ associated to the hypercharge group. Introducing the fields A_μ and Z_μ such that

$$\begin{aligned} B^\mu &= \cos\theta_{_W}\, A^\mu - \sin\theta_{_W}\, Z^\mu \\ W_3^\mu &= \sin\theta_{_W}\, A^\mu + \cos\theta_{_W}\, Z^\mu, \end{aligned} \tag{43}$$

with $\theta_{_W}$ an adjustable parameter, we re-express the relevant pieces of the lagrangian. Writing out explicitely the result for the quark sector we find

$$\left.\begin{aligned} \mathcal{L} = & (\bar{u}_{_L}\bar{d}_{_L}) \begin{pmatrix} \frac{2}{3}A^\mu & 0 \\ 0 & -\frac{1}{3}A^\mu \end{pmatrix} e\, \gamma_\mu \begin{pmatrix} u_{_L} \\ d_{_L} \end{pmatrix} \\ & + (\bar{u}_{_R}\bar{d}_{_R}) \begin{pmatrix} \frac{2}{3}A^\mu & 0 \\ 0 & -\frac{1}{3}A^\mu \end{pmatrix} e\, \gamma_\mu \begin{pmatrix} u_{_R} \\ d_{_R} \end{pmatrix} \end{aligned}\right\} \text{photon coupling} \tag{44}$$

$$\text{Z coup.} \left\{ \begin{array}{l} +(\bar{u}_{_L}\bar{d}_{_L}) \begin{pmatrix} (\frac{1}{2} - \frac{2}{3}\sin^2\theta_{_W})Z^\mu & 0 \\ 0 & (-\frac{1}{2} + \frac{1}{3}\sin^2\theta_{_W})Z^\mu \end{pmatrix} \frac{e}{\sin\theta_{_W}\cos\theta_{_W}}\ \gamma_\mu \begin{pmatrix} u_{_L} \\ d_{_L} \end{pmatrix} \\ \\ +(\bar{u}_{_R}\bar{d}_{_R}) \begin{pmatrix} (-\frac{2}{3}\sin^2\theta_{_W})Z^\mu & 0 \\ 0 & (\frac{1}{3}\sin^2\theta_{_W})Z^\mu \end{pmatrix} \frac{e}{\sin\theta_{_W}\cos\theta_{_W}}\ \gamma_\mu \begin{pmatrix} u_{_R} \\ d_{_R} \end{pmatrix} \end{array} \right. \tag{45}$$

where the parameters $g, g', \theta_{_W}$ are chosen such that:

$$g\sin\theta_{_W} = g'\cos\theta_{_W} = e \tag{46}$$

One finds (by construction!) that the A_μ field has the same couplings to the two helicity states of a given fermion, therefore it couples vectorially: we interpret this field as the photon which couples to the quarks with charges $e_u = \frac{2}{3}$ and $e_d = -\frac{1}{3}$. The other neutral field Z_μ, on the other hand, has both vector and axial-vector couplings to the fermion. The above equation can be written extremely simply as

$$\mathcal{L} = e\sum_{q=u,d} e_q\ \bar{q}\ \not{A}\ q\ +\ \frac{e}{\sin\theta_{_W}\cos\theta_{_W}}\sum_{q=u,d}\bar{q}\ \not{Z}\ (a_q + b_q\gamma_5)q \tag{47}$$

with

$$a_q = \frac{t_3}{2} - e_q\sin^2\theta_{_W}, \qquad b_q = \frac{t_3}{2}. \tag{48}$$

A similar derivation can be carried out for the leptonic sector. These couplings are in agreement with those of the physical Z boson once the Weinberg angle $\theta_{_W}$ (in fact introduced by Glashow!) is taken from experiment to be

$$\sin^2\theta_{_W} \sim .2322\ . \tag{49}$$

Considering what has been achieved until now, one finds that the model based on the $SU(2)_L\otimes U(1)_Y$ symmetry contains four gauge bosons: two charged ones with $(V-A)$ couplings to fermions and two neutral ones with couplings such that these bosons can be interpreted as the photon and the Z boson. The "only" difference with the real world is that in the present state of development of the model the gauge bosons are massless, because of the assumed exact gauge invariance and the fermions are also massless because of the left-right asymmetry of the gauge group. Counting the bosonic degrees of freedom of the model one realizes that three degrees of freedom are "missing", associated to the longitudinal polarisation states of the heavy vector bosons as summarized in the table.

	Model			Real World	
	degrees of freedom			degrees of freedom	
	transverse	longitudinal		transverse	longitudinal
W^-	2	0	W^-	2	1
W^+	2	0	W^+	2	1
Z	2	0	Z	2	1
γ	2	0	γ	2	0

In order to complete the model one should therefore introduce at least three new fields in the lagrangian. This will be done through a multiplet of scalar fields and it will be seen that, by the mechanism of spontaneous symmetry breaking of local gauge invariance, some of the scalar fields become the longitudinal polarization states and correlatively the vector bosons acquire a mass. The following argument may help understand the connection between a scalar field and a longitudinal polarisation state. Consider for example, the $q\bar{q}Z$ vertex for a boson in a longitudinal state of momentum q. It has the form

$$g\,[\bar{u}\gamma_\mu(a+b\gamma_5)u]\,\epsilon^\mu_{_L}(q) \sim g\,[\bar{u}\gamma_\mu(a+\ b\gamma_5)u]\,\frac{q^\mu}{M_{_Z}} \qquad |q^\mu| \gg M_{_Z} \tag{50}$$

This can be also interpreted, at asymptotic energies, as the derivative coupling of a scalar field ω with the fermionic current since, in momentum space, the derivative brings down a q^μ factor

$$g_\omega\,[\bar{u}\gamma_\mu(a+b\gamma_5)u]\,\partial^\mu\omega \rightsquigarrow g_\omega\,[\bar{u}\gamma_\mu(a+b\gamma_5)u]\,q^\mu\omega. \tag{51}$$

where the coupling g_ω necessarily has the dimension of the inverse of a mass. In the asymptotic limit a longitudinal boson couples to the fermions like a scalar with coupling $g_\omega \sim g/M_{_Z}$.

4 Spontaneous symmetry breaking

We proceed in steps and discuss, first, the case of a global symmetry and state the Golstone theorem. It is then applied specifically to the electro-weak model. Finally we make the symmetry local which leads us to the famous Brout-Englert-Higgs-Kibble mechanism.

4.1 Global symmetry breaking

Consider the very simple case of a complex scalar field

$$\varphi = \frac{1}{\sqrt{2}}(\varphi_1 + i\varphi_2) \tag{52}$$

which has two degrees of freedom $\varphi_1(x), \varphi_2(x)$. The lagrangian is invariant under the global phase transformation $\varphi(x) \to e^{i\alpha}\varphi(x)$ (α is constant) where

$$\mathcal{L} = \partial_\mu\varphi^*\partial^\mu\phi - V(\varphi) \text{ with the potential } V(\varphi) = -\mu^2|\varphi|^2 + h|\varphi|^4. \tag{53}$$

We assume μ^2 is negative so that the potential takes the well-known "Mexican hat" or "cul-de-bouteille" shape (depending on your cultural background!). The hamiltonian is

$$\begin{aligned} H &= \pi\ \partial_0\ \varphi - \mathcal{L}, \ \text{ with } \ \pi = \frac{\delta\mathcal{L}}{\delta\partial_0\varphi} = \partial_0\varphi^* \\ &= \underbrace{\partial_0\varphi^*\partial_0\varphi + |\vec{\nabla}\varphi|^2}_{H_{\text{kinetic}}} + V(\varphi). \end{aligned} \tag{54}$$

The (positive) kinetic part vanishes for static configurations and the full hamiltonian is minimal for constant values of the field given by

$$|\varphi_0| = \frac{v}{\sqrt{2}} = \frac{\mu}{\sqrt{2h}} \tag{55}$$

which defines the so-called vacuum expectation value v of the field φ in terms of the parameters of the lagrangian. Indeed, the quantum theory should be constructed from the lowest energy classical state which, in this case, is characterised by having its norm constrained by the above equation: such a state is called the classical vacuum. One immediately notices that the vacuum is degenerate since the application of a gauge transformation (phase change) does not affect the norm of the state. However to construct the quantum theory one needs to choose a particular vacuum, by imposing, for example, the classical vacuum field to be real *i.e.*

$$\varphi_0 = \frac{v}{\sqrt{2}} \tag{56}$$

This obviously amounts to breaking the symmetry of the vacuum since φ_0 is no more invariant under a gauge transformation, but the dynamical laws are still unbroken because they are given by the gauge invariant lagrangian eq. (53). This is the basis of "spontaneous symmetry breaking" in contradistinction to "explicit symmetry breaking" where the lagrangian itself would loose gauge invariance. To study the theory, we translate the original field by its vacuum expectation value

$$\varphi = \frac{1}{\sqrt{2}}(v + \varphi_1(x) + i\varphi_2(x)) \tag{57}$$

and, neglecting constant terms, the lagrangian becomes

$$\begin{aligned} \mathcal{L} \;\; = \;\; & \frac{1}{2}(\partial_\mu\varphi_1)^2 + \frac{1}{2}(\partial_\mu\varphi_2)^2 \;\; - \;\; hv^2\varphi_1^2 \\ & + hv\varphi_1(\varphi_1^2 + \varphi_2^2) \;\; + \;\; \frac{h}{4}(\varphi_1^2 + \varphi_2^2)^2 \end{aligned} \tag{58}$$

After spontaneous symmetry breaking, we are left with a model of two interacting real fields φ_1 and φ_2. The free theory is given by the first line of the equation above which shows that φ_2 is massless while φ_1 has a mass $m_{\varphi_1} = \sqrt{2hv^2}$. The interaction part is all contained in the second line of eq. (58) and there are cubic and quartic interactions between φ_1 and φ_2. Since, the initial lagrangian contained only two parameters, there are necessarily relations between the three parameters m_{φ_1}, the cubic coupling $g_{_3}$ and the quartic coupling $g_{_4}$ *e.g.*

$$g_{_3} = 2m^2_{\varphi_1}\; g_{_4}. \tag{59}$$

These features are a simple illustration of very general properties of spontaneous breaking of larger (non-abelian) group symmetry. In particular one can state the Goldstone theorem:

When a global symmetry is spontaneously broken there appear as many massless scalar modes (called the Goldstone bosons) as there are broken degrees of symmetry.

A proof of this theorem is now sketched. Consider a collection of n scalar fields $\varphi_i,\ i = 1, \cdots, n$ written as a column vector so that

$$\varphi^{^T} = (\varphi_{_1},\ \cdots, \varphi_{_n}) \tag{60}$$

The lagrangian density is formally written as

$$\mathcal{L} = \mathcal{L}(\varphi, \partial_\mu \varphi)_{\rm kin}\ -\ V(\varphi). \tag{61}$$

The vacuum of the model is defined by the conditions

$$\frac{\delta V}{\delta \varphi_i} = 0,\ \rightsquigarrow \text{vacuum:}\ \ \varphi^{0^T} = (\varphi^0_{_1},\ \cdots, \varphi^0_n) \tag{62}$$

One perturbs around the vacuum state

$$\varphi = \varphi^0 + \varphi',\quad i.e.\ \ \varphi_i = \varphi^0_i + \varphi'_i \tag{63}$$

so that the lagrangian (neglecting constant terms) is re-written

$$\mathcal{L} = \mathcal{L}(\varphi', \partial_\mu \varphi')_{\rm kin}\ - \frac{1}{2}\sum_{ij} \frac{\delta V}{\delta\varphi_i \delta\varphi_j}\varphi'_i\varphi'_j\ \oplus\ (\varphi'^3)\ \oplus\ (\varphi'^4) \tag{64}$$

where it is not necessary for our present purposes to specify the cubic nor the quartic couplings. By construction, there are no terms linear in the fields because we are expanding around the minimum of the potential. The quantity of interest is the quadratic term which defines the mass matrix

$$m^2_{ij} = \frac{\delta V}{\delta\varphi_i \delta\varphi_j}. \tag{65}$$

Consider now the action of an infinitesimal global gauge transformation. Its action on the fields is

$$\delta\varphi = i\ \alpha^J\ T^J\ \varphi, \qquad J = 1, \cdots, N, \tag{66}$$

where the T^J are the N generators ($n \times n$ matrices) of the group and the α^J are the N associated arbitrary parameters. If for some field configuration φ we have for a particular generator T^J,

$$T^J\varphi = 0,\ \ \rightsquigarrow\ \ \delta\varphi = i\alpha^J\ T^J\varphi = 0, \tag{67}$$

then we say that the configuration φ is invariant under the sub-group generated by T^J: the corresponding symmetry is unbroken. If, on the contrary, $T^J\varphi \neq 0$ the corresponding degree of symmetry is said to be spontaneously broken. Let us suppose now that the vacuum satisfies

$$\begin{aligned} &T^J\varphi_0 \neq 0 \qquad &&\text{for } J = 1, \cdots, N' \\ &T^J\varphi_0 = 0 \qquad &&\text{for } J = N'+1, \cdots, N, \end{aligned} \tag{68}$$

i.e. that the vacuum state breaks N' degrees of symmetry. The invariance of the lagrangian under the gauge transformation

$$\delta\varphi_i = i\ \alpha^J\ T^J_{ik}\ \varphi_k \tag{69}$$

for arbitrary α^J yields $\frac{\delta V}{\delta\varphi_i}\delta\varphi_i \equiv 0$. Taking the derivative of this relation it comes out

$$\frac{\delta^2 V}{\delta\varphi_j\delta\varphi_i}T^J_{il}\varphi_l \;+\; \frac{\delta V}{\delta\varphi_i}\delta_{ij} = 0 \;\rightsquigarrow\; m^2_{ji}T^J_{il}\;\varphi^0_l = 0. \tag{70}$$

Since this relation is automatically satisfied for $J = N'+1,\cdots,N$ one concludes that the mass matrix must have N' vanishing eigenvalues. Thus, N' fields φ'_i will be massless which are the Golstone bosons associated to the N' degrees of broken symmetry (*qed*).

4.2 Application to the electro-weak gauge group

In this case, the generators of the symmetry group will be $T^J = (\tau_1,\tau_2,\tau_3,Y)$, *i.e.* the generators of the weak isospin group and of the hypercharge. We introduce the complex scalar field, which is a doublet of $SU(2)$ ($t_{_\Phi} = \frac{1}{2}$),

$$\Phi = \frac{1}{\sqrt{2}}\begin{pmatrix} \varphi_{_1} \;-\; i\;\varphi_{_2} \\ \varphi_{_3} \;-\; i\;\varphi_{_4} \end{pmatrix} \tag{71}$$

and the standard scalar lagrangian

$$\mathcal{L}_S = \partial_\mu\Phi^\dagger\partial^\mu\Phi - V(\Phi),\;\; V(\Phi) = -\mu^2\Phi^\dagger\Phi \;+\; h\;(\Phi^\dagger\Phi)^2. \tag{72}$$

We will define the physical vacuum by

$$\Phi_0 = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \quad \text{with} \;\; v^2 = \frac{\mu^2}{h} \tag{73}$$

such that, of course, $\delta V/\delta\Phi|_{\Phi=\Phi_0} = 0$. Since we require the electric charge to be conserved after symmetry breaking, we have to enforce, following the previous reasoning, that the charge generator acting on the vacuum state should vanish. Following the Gell-Mann/Nishijima relation eq. (35) we need

$$Q\;\Phi_0 = \frac{1}{2}\;(\tau_3 + Y)\;\Phi_0 = \begin{pmatrix} \frac{1}{2}+\frac{y}{2} & 0 \\ 0 & -\frac{1}{2}+\frac{y}{2} \end{pmatrix}\;\begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \;=\; 0 \tag{74}$$

implying that the hypercharge of the scalar field must be $y_{_\Phi} = 1$ to ensure charge conservation in the broken theory. As in the abelian case, we can study the system around the classical minimum and expand the scalar field around its vacuum expectation value v

$$\Phi = \begin{pmatrix} \frac{-i}{\sqrt{2}}(\omega_{_1}(x) - i\omega_{_2}(x)) \\ \frac{1}{\sqrt{2}}(v + H(x) - i\omega_{_3}(x)) \end{pmatrix} \;=\; \begin{pmatrix} \omega^\dagger(x) \\ \frac{1}{\sqrt{2}}(v + H(x) - i\omega_{_3}(x)) \end{pmatrix}, \tag{75}$$

so that the scalar potential $V(\Phi)$ becomes

$$V(\Phi) = hv^2H^2 \;+\; hv\;H(H^2+\vec{\omega}^2) \;+\; \frac{h}{4}\;(H^2+\vec{\omega}^2)^2 \tag{76}$$

showing that the triplet of ω_i fields (one neutral and two charged fields) is massless while the neutral H field acquires a mass

$$M_{_H} = \sqrt{2hv^2}. \tag{77}$$

All these fields are coupled together with a strength which can be read off the equation above.

4.3 Local symmetry breaking and the Brout-Englert-Higgs-Kibble mechanism

Armed with this lengthy preliminaries we now go directly into the spontaneous breaking of the local gauge symmetry $SU(2)_L \otimes U(1)_Y$ down to $U(1)_{emg}$. Let us state the results before diving into an ocean of technicalities. The case of a global symmetry has just been analysed and led to the appearence of three massless (Goldstone) bosons and a massive one. When the symmetry is made local these massless bosons turn out to be unphysical, in the sense that they can be gotten rid off by a gauge transformation, but instead, three gauge bosons (a neutral one and the two charged ones) become massive and therefore acquire longitudinal polarisation states which are the Goldstone modes in disguise.

To implement symmetry breaking we first have to extend the electro-weak lagrangian eq. (38) to include the scalar field contribution $\mathcal{L}_S$ as well as the interaction of the new scalar field with the fermions $\mathcal{L}_Y$ (where Y stands for Yukawa) so that the electro-weak lagrangian is

$$\mathcal{L} \;=\; \mathcal{L}_F \;+\; \mathcal{L}_G \;+\; \mathcal{L}_S \;+\; \mathcal{L}_Y. \tag{78}$$

4.3.1 The scalar lagrangian $\mathcal{L}_S$

We concentrate for the moment on $\mathcal{L}_S$ which drives the spontaneous breaking of the local electro-weak symmetry. No other fields such as fermions or gauge bosons can acquire vacuum expectation values otherwise the physical vacuum would have some angular momentum or other non-vanishing quantum numbers. The scalar lagrangian has the form

$$\mathcal{L}_S = |(\partial^\mu - i\; g\; \frac{\vec{\tau}}{2} \cdot \vec{W}^\mu - i\; \frac{1}{2}\; g'\; B^\mu)\Phi|^2 \;-\; V(\Phi) \tag{79}$$

where making the symmetry local amounts to replacing the ordinary derivative in eq. (72) by the appropriate covariant derivative acting on a **2** field of a local $SU(2)$ symmetry which also has non-zero hypercharge (see eq. (39)). To analyse the effects of symmetry breaking we substitute into this equation the vacuum expectation value Φ_0 of Φ (eq. (73)) and also work with the "physical" A_μ and Z_μ fields of eq. (43) rather than with $\vec{W}_\mu$ and B_μ. We have

$$\begin{aligned}
\mathcal{L}_S|_{\Phi=\Phi_0} \;&=\; \Bigg| -\frac{ie}{\sqrt{2}\sin\theta_{_W}} \begin{pmatrix} 0 & W^-_\mu \\ W^+_\mu & 0 \end{pmatrix} \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \\
&\quad -ie \begin{pmatrix} A_\mu - \frac{\cos^2\theta_{_W} - \sin^2\theta_{_W}}{2\sin\theta_{_W}\cos\theta_{_W}} Z_\mu & 0 \\ 0 & -\frac{1}{2\sin\theta_{_W}\cos\theta_{_W}} Z_\mu \end{pmatrix} \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \Bigg|^2 - V(\Phi_0) \\
&=\; \frac{e^2}{4\sin^2\theta_{_W}} v^2 W^+_\mu W^{-\mu} \;+\; \frac{e^2}{8\sin^2\theta_{_W}\cos^2\theta_{_W}} v^2 Z_\mu Z^\mu \;-\; V(\Phi_0)
\end{aligned} \tag{80}$$

This equation tells us immediately that the charged bosons $W^\pm$ acquire the mass $M^2_{_W} = \frac{e^2}{4\sin^2\theta_{_W}} v^2$, the Z boson a mass $M^2_{_Z} = \frac{e^2}{4\sin^2\theta_{_W}\cos^2\theta_{_W}} v^2$, while the photon remains massless as no quadratic term in A_μ appears in the lagrangian. Note the important relation

$$M_{_W} = M_{_Z} \cos\theta_{_W} \tag{81}$$

which is a prediction of the model once the boson couplings to fermions have been determined from experiment. The vanishing of the photon mass is a consequence of the surviving exact gauge symmetry $U(1)_{emg}$. We have the relation $v = \frac{\sin\theta_{_W} M_{_W}}{\sqrt{\pi\alpha}}$ between the vacuum expectation value of the scalar field and the physical parameters and, plugging in numerical values, we find $v \sim 250$ GeV, which is the basis for the claim, made in the introduction, that the non-abelian symmetry is broken at the scale of 250 GeV.

Turning now to the study of oscillations around the vacuum Φ_0 we write the scalar Φ in the form eq. (75). One immediately notices that the three fields $\omega_i(x)$ can be gauged away since we can always choose the parameters $\vec{\alpha}(x)$ of a local gauge transformation such as

$$\exp(-i\frac{\vec{\tau}}{2}\vec{\alpha}(x))\ \Phi\ =\ \begin{pmatrix} 0 \\ \frac{v+H(x)}{\sqrt{2}} \end{pmatrix}, \tag{82}$$

showing that the fields $\omega_i(x)$ can be removed from the lagrangian altogether and therefore are not physical. This is consistent and, in fact, it must be so since the massive vector bosons have now a mass and therefore an extra degree of (longitudinal) polarisation so that the total number of degrees of freedom is not changed by the breaking of the symmetry. Of course, explicit gauge invariance will be lost since a particular gauge (the so-called unitary gauge) has been chosen. Parameterising the scalar field as in the right-hand side of eq. (82) (*i.e.* keeping only the physical degrees of freedom) one immediately reads off from the lagrangian $\mathcal{L}_S$ the mass and couplings of the famed Higgs field H. From $V(\Phi)$ (see eq. (76) setting $\vec{\omega} = 0$) one gets the mass

$$M_{_H} = \sqrt{2h}\ v \tag{83}$$

and the triple and quadruple self-couplings respectively

$$h_{_3} \sim hv \sim \frac{M_{_H}}{\sqrt{h}} \sim e\frac{M_{_H}^2}{M_{_W}},\quad \text{and}\quad h_{_4} \sim h \sim e^2\frac{M_{_H}^2}{M_{_W}^2}. \tag{84}$$

while from the kinetic part of $\mathcal{L}_S$ come out the couplings of the Higgs particle to the gauge bosons. To derive them it is enough to make the substitution $v \to v + H$ in eq. (80) and one obtains

− the triple coupling (Higgs-W^+-W^-) $= \frac{e^2}{2\sin^2\theta_{_W}} v = \frac{e}{\sin\theta_{_W}} M_{_W}$

− the quadruple coupling (Higgs-Higgs-W^+-W^-) $= \frac{e^2}{2\sin^2\theta_{_W}}$.

The derivation of similar couplings to the Z boson is left as an exercise.

As discussed before, the existence of the massive gauge bosons leads to a bad behaviour of the amplitudes at high energies because of the longitudinal polarisation states. It can be explicitely verified that, at the tree level, those divergences are cancelled when keeping all diagrams including those with the Higgs particle. At higher orders, loop diagrams may involve the massive gauge boson propagator which is of the form

$$\Delta^{\mu\nu} = i(g^{\mu\nu} - \frac{q^\mu q^\nu}{M^2})/(q^2 - M^2) \tag{85}$$

which does not converge to 0 when $q^\mu \to \infty$ because of the $q^\mu q^\nu/M^2$ term: this leads to an apparently non-renormalisable theory. As explained in detail for the abelian case in Piguet's

lectures [8], the way out is not to work in the unitary gauge which keeps only physical degrees of freedom, but rather to work in a "renormalisable" gauge (for example, in the 't Hooft gauge) where the Golstone modes $\vec{\omega}$ are explicitely kept in the calculation. It can be proven (equivalence theorem) that these modes are related to the longitudinal polarisation states of the gauge bosons.

4.3.2 The Yukawa lagrangian $\mathcal{L}_Y$ and fermion masses and couplings

The scalar field Φ can couple to fermions. The requirement for such couplings to exist is that the corresponding terms in the Lagrangian be invariant under a $SU(2)_L \times U(1)_Y$ transformation (before the spontaneous breaking of this symmetry is implemented of course!). Let us recall that ψ_{e_L} and ψ_{q_L} of eq. (25) and Φ of eq. (71) are **2** under $SU(2)$ *i.e.* they transform as

$$\delta\psi = i\ \frac{\vec{\tau}}{2}\ \vec{\alpha}\ \psi,\ \cdots\ , \delta\bar{\psi} = -i\ \bar{\psi}\ \frac{\vec{\tau}}{2}\ \vec{\alpha},\ \cdots \tag{86}$$

so that $\bar{\psi}_{e_L}\Phi$, $\bar{\psi}_{q_L}\Phi$ are invariant under a $SU(2)_L$ transformation. One easily shows that the combination $i\tau_2\Phi^*$ is also a **2** and consequently the quantites $\bar{\psi}_{e_L}(i\tau_2\Phi^*)$ and $\bar{\psi}_{q_L}(i\tau_2\Phi^*)$ are also invariant. Considering now the transformation properties under $U(1)_Y$ it turns out that the terms $\bar{\psi}_{e_L}\Phi\ e_R$, $\bar{\psi}_{q_L}\Phi\ d_R$, $\bar{\psi}_{q_L}(i\tau_2\Phi^*)\ u_R$ are all invariant: this is easily seen using the fermion hypercharge assignements given in table (37) and the fact that $y_\Phi = -y_{\Phi^*} = 1$ (eq. (74)). We do not include a term with ν_R as we assume the right-handed neutrino does not couple to the physical world in agreement with experiment. The Yukawa lagrangian then takes the form

$$\mathcal{L}_Y = c_d\ \bar{\psi}_{q_L}\Phi d_R + c_u\ \bar{\psi}_{q_L}(i\tau_2\Phi^*)u_R + c_e\ \bar{\psi}_{e_L}\Phi e_R + \text{h.c.} + \text{other families}, \tag{87}$$

where we have explicitely written out the terms involving the first family of fermions $(\nu, e; u, d)$. Six other parameters should be similarly introduced for the couplings of the second and third families so that nine new parameters appear in the model. Implementing spontaneous symmetry breaking, in the unitary gauge, *i.e.* substituting in $\mathcal{L}_Y$ the expression of Φ as given in the right-hand side of eq. (82), we derive

$$\mathcal{L}_Y = c_d\ \frac{v+H}{\sqrt{2}}\bar{d}d\ +\ c_u\ \frac{v+H}{\sqrt{2}}\bar{u}u\ +\ c_e\ \frac{v+H}{\sqrt{2}}\bar{e}e\ +\ \text{other families}. \tag{88}$$

From this expression we relate the mass of a fermion f to the vacuum expectation value v *via*

$$M_f = c_f\frac{v}{\sqrt{2}}. \tag{89}$$

This is not a prediction of the theory since the parameters c_f are unknown and will be adjusted so as to obtain the "physical" mass of the corresponding fermion. Furthermore, no relation is expected between the masses of partners of a given family since one parameter is introduced for each of the fermion type in a family. One may remark that the only "prediction" is that the neutrino remains massless as a consequence of the absence a right-handed neutrino. On

the other hand the Higgs couplings to the fermions are predicted, if the fermion masses are known,

$$g_f = \frac{c_f}{\sqrt{2}} = \frac{e}{2\sin\theta_{_W}}\,\frac{M_f}{M_{_W}}, \tag{90}$$

where eq. (89) and the relation after eq. (80) have been used: the Higgs particle couples to a fermion flavour in proportion to its mass, implying that the top quark could play a major role in the production and/or decay of the Higgs particle ($M_t \sim 175$ GeV) while the electron contribution can be safely neglected ($M_e = .511\ 10^{-3}$ GeV). As an application we consider



Figure 2: *Higgs production mechanism at hadron-hadron colliders. The dominant contribution arises from a top quark loop.*

Higgs production in proton-proton colliders such as LHC (at a center of mass energy $\sqrt{s} = 14$ TeV). For a Higgs mass below about 500 GeV the dominant process is gluon-gluon fusion where the effective Higgs coupling to the gluon-gluon system is *via* a quark loop as indicated in the figure. Because of the very large top mass the loop contribution is dominated by the top quark. The possible direct $t\bar{t} \to H$ is of course suppressed since the density of (virtual) $t, \bar{t}$ in the proton is negligible.

4.4 Family mixing and the Kobayashi-Maskawa matrix

The above discussion has been considerably simplified since it completely ignored mixing between generations. However it turns out that the charged weak interactions are not diagonal in flavour, or to put it more precisely, the flavour basis defined as the basis which diagonalises the fermion mass matrix is not the natural one to study charged current interactions. Furthermore there is no reason for the absence in the Yukawa lagrangian of terms which mix families. To study these features let us introduce a new basis of quark flavours *e.g.* ($u'_{_{Lj}}$) related to the usual one ($u_{_{Lj}}$) by

$$u_{_{Li}} = S^{u_L}_{ij}\, u'_{_{Lj}}, \qquad i,j = 1,2,3 \tag{91}$$

where the indices i, j run over the number of families. This can be written in a matrix form $\mathbf{u}_{_L} = \mathbf{S}^{u_L}\,\mathbf{u}'_{_L}$ and similarly for the right-handed *up* sector as well as the left-handed and right-handed *down* sectors. As will be seen later the matrices $\mathbf{S}^{u_L}, \mathbf{S}^{u_R}, \cdots$ are unitary. The quark

content of the $SU(2)_L \otimes U(1)_Y$ lagrangian is written in general (see eq. (38))

$$\mathcal{L}_F = \sum_i (\bar{u}'_{_{Li}} \; \bar{d}'_{_{Li}}) \; i \not{D}_{_L} \begin{pmatrix} u'_{_{Lj}} \\ d'_{_{Lj}} \end{pmatrix} \; + \; \sum_i \bar{u}'_{_{Ri}} \; i \not{D}_{_R} \; u'_{_{Ri}} \; + \; \sum_i \bar{d}'_{_{Ri}} \; i \not{D}_{_R} \; d'_{_{Ri}}, \tag{92}$$

while the most general Yukawa lagrangian takes the form after symmetry breaking (see eqs. (87),(88))

$$\begin{aligned} \mathcal{L}_Y &= \frac{v}{\sqrt{2}} \sum_{ij} (\bar{u}'_{_{Li}} c^u_{ij} u'_{_{Rj}} \; + \; \bar{d}'_{_{Li}} c^d_{ij} d'_{_{Rj}} \; + \; \text{h.c.}) \\ &= \frac{v}{\sqrt{2}} (\mathbf{u}'_{_L} \mathbf{C}_u \mathbf{u}'_{_R} \; + \; \mathbf{u}'_{_R} \mathbf{C}^\dagger_u \mathbf{u}'_{_L} \; + \; \mathbf{d}'_{_L} \mathbf{C}_d \mathbf{d}'_{_R} \; + \; \mathbf{d}'_{_R} \mathbf{C}^\dagger_d \mathbf{d}'_{_L}) \end{aligned} \tag{93}$$

where the complex, 3×3, $\mathbf{C}_u, \mathbf{C}_d$ matrices are the generalised Yukawa couplings. The most general such matrices can be written as a product

$$\mathbf{C}_u = \mathbf{M}_u \; \mathbf{T}_u, \qquad \mathbf{C}_d = \mathbf{M}_d \; \mathbf{T}_d \tag{94}$$

with $\mathbf{M}_u$ a hermitian matrix ($\mathbf{M}_u = \mathbf{M}^\dagger_u$) and $\mathbf{T}_u$ a unitary matrix ($\mathbf{T}^{-1}_u = \mathbf{T}^\dagger_u$). The hermitian matrix can be diagonalised by a unitary transformation, $\mathbf{M}_u = \mathbf{S}^{-1}_u \mathbf{m}_u \mathbf{S}_u$ where $\mathbf{m}_u$ is diagonal with real eigenvalues, and similarly for the *down* sector. The Yukawa lagrangian reduces to the very simple diagonal form

$$\begin{aligned} \mathcal{L}_Y &= \bar{\mathbf{u}}_{_L} \; \mathbf{m}_u \; \mathbf{u}_{_R} \; + \; \bar{\mathbf{u}}_{_R} \; \mathbf{m}_u \; \mathbf{u}_{_L} \; + \; \bar{\mathbf{d}}_{_L} \; \mathbf{m}_d \; \mathbf{d}_{_R} \; + \; \bar{\mathbf{d}}_{_R} \; \mathbf{m}_d \; \mathbf{d}_{_L} \\ &= \bar{\mathbf{u}} \; \mathbf{m}_u \; \mathbf{u} \; + \; \bar{\mathbf{d}} \; \mathbf{m}_d \; \mathbf{d} \end{aligned} \tag{95}$$

when written in terms of the physical basis related to the original one by

$$\begin{aligned} \mathbf{u}_{_L} &= \mathbf{S}_u \; \mathbf{u}'_{_L}, \qquad \mathbf{u}_{_R} = \mathbf{S}_u \; \mathbf{T}_u \; \mathbf{u}'_{_R} \\ \mathbf{d}_{_L} &= \mathbf{S}_d \; \mathbf{d}'_{_L}, \qquad \mathbf{d}_{_R} = \mathbf{S}_d \; \mathbf{T}_d \; \mathbf{d}'_{_R}. \end{aligned} \tag{96}$$

We remark that, as advertised before, the transformation from the primed basis to the unprimed one is unitary since such are the $\mathbf{S}, \mathbf{T}$ matrices. The components of $\mathbf{u}$ in which the mass matrix is diagonal are, by definition, the "physical" quark fields (u, c, t) of definite mass eigenstate (*idem* for the d sector). Having achieved a simple form for the Yukawa lagrangian we re-write now the gauge part $\mathcal{L}_F$ in terms of these physical fields. Singling out the neutral current interactions we have

$$\mathcal{L}_F(\text{neutral current}) = \sum_i (\bar{u}'_{_{Li}} \; i \not{D}_{_L} \; u'_{_{Li}} \; + \; \bar{u}'_{_{Ri}} \; i \not{D}_{_R} \; u'_{_{Ri}}) \; + \; d' \text{ sector}, \tag{97}$$

in which we keep only the diagonal (in $SU(2)$) part of the operator $\not{D}_{_L}$. Because of the unitarity of the transformations within the left-handed bases and the right-handed bases the above lagrangian immediately reduces itself to

$$\begin{aligned} \mathcal{L}_F(\text{neutral current}) &= \bar{\mathbf{u}}_{_L} \; i \not{D}_{_L} \; \mathbf{u}_{_L} \; + \; \bar{\mathbf{u}}_{_R} \; i \not{D}_{_R} \; \mathbf{u}_{_R} \\ &+ \; \bar{\mathbf{d}}_{_L} \; i \not{D}_{_L} \; \mathbf{d}_{_L} \; + \; \bar{\mathbf{u}}_{_R} \; i \not{D}_{_R} \; \mathbf{u}_{_R} \end{aligned} \tag{98}$$

This equation is the basis for the slogan that, in the Standard Model and in agreement with experiments, there is "no flavour-changing neutral current", in other words the neutral current is diagonal in flavour space and does induce transition between the c quark and the u quark or between the b quark and the d quark for example. The case of the charged current pieces is more involved because it couples the *up* sector and the *down* sector which do not transform with the same unitary matrices and, as a consequence, there is no reason for the charged current interactions to be diagonal in the basis which diagonalises the mass matrix. Indeed we have, from eqs. (92, 39)

$$\begin{aligned} \mathcal{L}_F(\text{charged current}) &= g\ \bar{\mathbf{u}}'_{_L}\ \gamma^\mu\ \mathbf{d}'_{_L}\ W^-_\mu\ +\ \text{h.c.} \\ &= g\ \bar{\mathbf{u}}_{_L}\ \gamma^\mu\ \mathbf{S}_u\mathbf{S}_d^\dagger\ \mathbf{d}_{_L}\ W^-_\mu\ +\ \text{h.c.} \end{aligned} \tag{99}$$

where the last equation is obtained from eq. (96). Obviously the matrix $\mathbf{V} = \mathbf{S}_u\mathbf{S}_d^\dagger$ is unitary and it depends therefore on 3^2 real parameters. Since $(2\times 3-1)$ parameters can be absorbed in a redefinition of the phases of the six fermion fields (one overall phase cannot be gotten rid of) there remain four independent parameters: these can be chosen as three real angles and a phase and it is convenient to write in the approximate Wolfenstein parameterisation:

$$\mathbf{V} = \begin{pmatrix} 1-\lambda^2/2 & \lambda & A\lambda^3(\rho-i\eta) \\ -\lambda & 1-\lambda^2/2 & A\lambda^2 \\ A\lambda^3(1-\rho-i\eta) & -A\lambda^2 & 1 \end{pmatrix}. \tag{100}$$

This is the famous Kobayashi-Maskawa (KM) matrix which generalises to three families the Cabibbo angle, $\sin\theta_{_C} = \lambda \sim .22$ introduced long ago to deal with the mixing of two families. This matrix shows that the charged current transition, for example, of a d quark to u, c, t quarks takes place with amplitudes which are proportional to $(1-\lambda^2/2), \lambda, A\lambda^3(\rho-i\eta)$ respectively. The calculation of processes which are used by experimentalists to measure accurately the various parameters of the KM matrix and to check its unitarity (relations such as $V^*_{ud}V_{ub} + V^*_{cd}V_{cb} + V^*_{td}V_{tb} = 0$ where the V_{ij} are the KM matrix elements) is a very active area of particle physics phenomenology at present. The phase factor η is responsible for CP violation in the Standard Model. The measurement of this CP violating parameter, in kaon and B meson systems, for example, is of great theoretical interest in order to understand the origin of CP violation and of great practical importance since it may be related to the origin of the baryon asymmetry in the universe.

The absence of a KM mixing matrix for the leptonic sector requires a comment. We assume (and it is supported by experiments) that the right-handed neutrinos decouple from the observed world. As a consequence, as mentioned above, the neutrinos ν_e, ν_μ, ν_τ remain massless even after spontaneous symmetry breaking and there is no mass matrix where the "physical" neutrino states can be defined. When studying the weak-current transition from charged leptons to neutrinos we can introduce a matrix $\mathbf{V}_\nu$ similar to the KM matrix of eq. (100) for the quark sector. We are free now to define the physical neutrino states by

$$\mathbf{V}_\nu^\dagger\ \boldsymbol{\nu} = \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}$$

i.e. to define the physical states as those for which the charged weak current is diagonal in lepton flavour.

5 Conclusions

Putting all together, the lagrangian which contains the dynamics of particle physics, as described by the Standard Model, is decomposed into

$$\mathcal{L} = \mathcal{L}_{QCD} \; + \; \mathcal{L}_G \; + \; \mathcal{L}_F \; + \; \mathcal{L}_S \; + \; \mathcal{L}_Y$$

where each piece has been previously defined. If one attempts to count the number of parameters introduced we arrive at:

- $SU(3)$ (QCD) gauge invariance: 1 coupling
- $SU(2) \otimes U(1)$ gauge invariance: 2 couplings + the Weinberg angle $\theta_{_W}$
- spontaneous symmetry breaking from $\mathcal{L}_S$: 2 parameters
- Yukawa couplings in $\mathcal{L}_Y$: 9 couplings + 4 KM parameters,

so one already has, at least, 19 parameters which is not a satisfactory situation for a minimal model! This has prompted an intensive continuing search for higher hidden symmetries such as supersymmetry, for example, and (a) Minimal Supersymmetry Standard Model(s) have (has) been constructed [9], [10]. The very unfortunate situation is that in such models the number of fields is more than doubled compared to the Standard Model as no supersymmetric multiplets can be filled with only known particles. Furthermore, while in principle supersymmetry breaking can in turn trigger electroweak symmetry breaking, one does not know yet how supersymmetry is dynamically broken. Thus one is lead to describe it in a effective way which requires many more parameters than in the Standard Model. It seems that, at present, the remedy is worse than the disease but there are still hopes .....

For references see *e.g.* [3, 4, 11, 12]

Acknowledgments

I am grateful to F. Boudjema, J.Ph. Guillet and E. Pilon for discussions and for a careful reading of this text. I also thank F. Gieres, M. Kibler, C. Lucchesi and O. Piguet, the organisers of the Vth Séminaire Rhôdanien de Physique at Dolomieu for a very enjoyable and educational meeting.

References

[1] *An introduction to quarks and partons*, F.E. Close, Academic Press (1979), London.

[2] *Quarks and leptons*, F. Halzen, A.D. Martin, John Wiley & sons.

[3] *Gauge theories of the strong, weak and electromagnetic interactions*, C. Quigg, The Benjamin/Cummings Publishing Company (1983), London.

[4] *Gauge theories of elementary particles*, T.P. Cheng and L.F. Li, Oxford University Press (1983).

[5] *Foundations of Quantum Chromodynamics*, T. Muta, World Scientific (1987), Singapore.

[6] *Quantum gauge theories*, S. Pokorski, Cambridge University Press (1987).

[7] *QCD and collider physics*, R.K. Ellis, W.J. Stirling, B.R. Webber, Cambridge University Press (1996).

[8] O. Piguet, *Proceedings of the Vth Séminaire Rhôdanien de Physique*, F. Gieres, M. Kibler, C. Lucchesi, O. Piguet *eds.*, Editions Frontières, 1997, Paris.

[9] J.P. Derendinger, see [8].

[10] H.P. Niles, see [8].

[11] *Quantum field theory*, L.H. Ryder, Cambridge University Press (1996).

[12] *An introduction to quantum field theory*, M.E. Peskin, D.V. Schröder, Addison-Wesley (1995), Reading, USA.